

Egg clutch structure of *Rhinella rumbolli* (Anura: Bufonidae), a toad from the Yungas of Argentina, with a review of the reproductive diversity in *Rhinella*

MARTÍN O. PEREYRA¹, M. FLORENCIA VERA CANDIOTI², JULIÁN FAIVOVICH^{1,3} & DIEGO BALDO⁴

¹) División Herpetología, Museo Argentino de Ciencias Naturales ‘Bernardino Rivadavia’ – CONICET, Ángel Gallardo 470, C1405DJR, Buenos Aires, Argentina

²) Instituto de Herpetología, Fundación Miguel Lillo – CONICET, Miguel Lillo 251, T4000JFE, San Miguel de Tucumán, Argentina

³) Departamento de Biodiversidad y Biología Experimental, Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires, Intendente Güiraldes 2160, C1428EGA, Buenos Aires, Argentina

⁴) Laboratorio de Genética Evolutiva, Instituto de Biología Subtropical (CONICET – UNaM), Facultad de Ciencias Exactas Químicas y Naturales, Universidad Nacional de Misiones, Félix de Azara 1552, N3300LQF, Posadas, Argentina

Corresponding author: MARTÍN O. PEREYRA, e-mail: mopereyra@gmail.com

Manuscript received: 31 July 2014

Accepted: 2 September 2014 by STEFAN LÖTTERS

Abstract. *Rhinella rumbolli* is a poorly known, medium-sized toad endemic to the Yungas of Argentina. Recent fieldwork allowed observing its peculiar oviposition mode, which is described in this paper. A review of literature and examination of museum material indicate that oviposition in *Rhinella* can vary from strings of eggs in a gelatinous tube (uniserial, biserial or multiserial) to open clumps. Clutch features in *Rhinella* provide informative variation for taxonomic and phylogenetic studies and help to understand the relationships in this large genus of true toads.

Key words. Argentina, *Rhinella veraguensis* group, oviposition modes, open clumps, strings, water mould.

Introduction

Rhinella is a species-rich bufonid genus, widely distributed throughout different Neotropical ecoregions (FROST 2014). Its 87 species are grouped into seven species groups: the *R. acrolopha* (formerly *Rhamphophryne*), *R. crucifer*, *R. granulosa*, *R. margaritifera*, *R. marina*, *R. spinulosa*, and *R. veraguensis* groups (DUELLMAN & SCHULTE 1992, PRAMUK 2006, GRANT & BOLIVAR-G. 2014), although the monophyly of some of them has not been properly corroborated (PRAMUK 2006, CHAPARRO et al. 2007, PYRON & WIENS 2011; MORAVEC et al. 2014). The reproductive biology of several species of the *R. marina*, *R. granulosa*, *R. crucifer*, *R. margaritifera*, and *R. spinulosa* groups is relatively well documented. However, this knowledge is deficient and limited merely to tadpole descriptions in some species of the *R. veraguensis* group, and to reports of number, colouration, and size of presumably mature oocytes in species of the *R. acrolopha* group (Appendix 1).

Rhinella rumbolli (CARRIZO, 1992) is a medium-sized toad from the Yungas of Argentina. This species is a largely aquatic inhabitant of forest streams and rivers, and males vocalize at the stream margins where the current is slower (CARRIZO 1992, HAAD et al. 2014). Adults have been ob-

served active or hiding in the leaf litter near rivers during the day (MOP pers. obs., D.E. CARDOZO & J.M. FERRO pers. comms.). The tadpoles of *R. rumbolli* have features intermediate between the morphology of pond tadpoles and that of gastromyzophorous forms exclusive to the *R. veraguensis* group (HAAD et al. 2014).

In this paper, we describe the clutch structure and reproductive mode of *Rhinella rumbolli*, and review the current knowledge on clutch structures and oviposition modes in *Rhinella*. In addition, we report the infection of clutches by water moulds of the family Saprolegniaceae. Finally, we interpret these findings in the context of the current phylogenetic hypotheses of *Rhinella* and discuss their systematic relevance.

Material and methods

A field survey was conducted on 25–26 August 2013, at Trigo Pampa River, Ocloyas (Jujuy, Argentina; 23°55'33.2" S, 65°15'13.1" W, datum WGS84; 1,500 m above sea level). Trigo Pampa is a typical mountain river, with bedrock and regimes of large floods during summer. The habitat corresponds to a type of rainforest known as ‘cloud forest’, and

is included in the Yungas Ecoregion (CABRERA 1994). Numerous males of *Rhinella rumbolli* were heard calling between 20:00 and 23:30 hours, and at this time several pairs were observed in axillar amplexus in a mountain river. A few specimens of *R. arenarum* and *Hypsiboas riojanus* were also active in the area. The next morning, egg clutches were collected from the same stream and preserved in 10% formalin, and many were retained for rearing the embryos. We assigned the clutches to *R. rumbolli* by rearing some of them to tadpoles, which can be easily identified (see HAAD et al. 2014), and confirmed their identity by comparison with a complete clutch (LGE 6415) that we obtained from an amplexant pair in the laboratory (LGE 6427 x LGE 6428). Eggs per clutch were counted and egg diameters measured to the nearest 0.01 mm using a Leica M205 A stereomicroscope with a DFC295 camera.

In order to explore variation among the species of *Rhinella*, published descriptions on reproductive biology (oviposition mode, number of eggs/oocytes, pigmentation, and ovum size) were compiled and are summarized in Appendix 1. This table also contains additional information about the reproductive biology of taxa with deficient data, which we obtained from photographic records, egg clutches in herpetological collections, and/or dissection of gravid females.

The terminology used to describe clutch structures is that of SALTHER (1963) and ALTIG & MCDIARMID (2007). For the composition of species groups of *Rhinella*, we follow DUELLMAN & SCHULTE (1992) with some modifications based on subsequent publications (e.g., BALDISERA et al. 2004, PRAMUK 2006, NARVAES & RODRIGUES 2009; GRANT & BOLIVAR-G. 2014). Studied specimens are housed in the following herpetological collections: División Herpetología, Museo Argentino de Ciencias Naturales 'Bernardino Rivadavia' – CONICET (MACN), Buenos Aires, Argentina; Instituto de Herpetología, Fundación Miguel Lillo (FML), Tucumán, Argentina; and Laboratorio de Genética Evolutiva (LGE), Instituto de Biología Subtropical (CONICET – Universidad Nacional de Misiones), Posadas, Misiones, Argentina. All the examined material is listed in Appendix 2.

Results

We found 25 clutches of *Rhinella rumbolli* in natural situations. All clutches had been deposited in places with shallow, slow-flowing or stagnant water, on the nude sandy bottom or among submerged vegetation of the river (mode




Figure 1. A, B) Clutches of *Rhinella rumbolli* photographed in situ in the Trigo Pampa river, at different stages of development (scale bars = 2 cm); the arrow points out hatched embryos at GOSNER (1960) stages 17–18; C) Detail of infected eggs (scale bar = 2 mm); D) showing the reproductive structures of water mould (scale bar = 0.5 mm).

2 of HADDAD & PRADO 2005). Most clutches were severely infected with water mould (Oomycota: Saprolegniaceae). In less severe cases, recently hatched embryos and successive developmental stages were observed near the spoiled eggs (Fig. 1).

A clutch of *Rhinella rumbolli* consists mostly of a mono-layered clump of eggs and lacks a surrounding matrix. It is an “open clump” as defined by ALTIG & MCDIARMID (2007) in that the surface of adjacent eggs is firm enough to form interstices (Fig. 2A). The only difference to ALTIG & MCDIARMID’s (2007) definition, where clumps do not have outlier eggs, is that we observed single and paired egg groups at the edges of the main group. The clutch obtained in the lab, the only one we are certain is complete, was approximately 20 cm in diameter, and consisted of 2,177 eggs.

The individual eggs are relatively large (3.85 mm in average diameter) and have a thick (about 0.97 mm), sticky jelly that makes adjacent eggs adhere to each other. The ova are also relatively large (about 1.91 mm) and have a pigmented, dark animal pole and a whitish vegetal pole.

The dissection of an adult female of *Rhinella quechua*, another species of the *R. veraguensis* group, failed to produce any oocytes from the ovisac, and we only found mature ovarian oocytes. The right ovary contained 507 oocytes with a pigmented animal pole and a mean diameter of 1.27 ± 0.05 mm (range 1.19–1.37; $N = 20$). Egg counts and descriptions from preserved clutches of *R. achalensis*, *R. arenarum*, *R. azarai*, *R. cf. cerradensis*, *R. fernandezae*, *R. major*, *R. ornata*, *R. quechua*, *R. schneideri*, and *R. spinulosa* are detailed in Appendix 1.




Figure 2. Clutches from nine species of the *Rhinella veraguensis* (RV), *R. granulosa* (RG), *R. marina* (RM), *R. crucifer* (RC), and *R. spinulosa* (RS) groups. The insets show details of each clutch and the arrangement of eggs. A) *R. rumbolli* RV (LGE 6415), portion of a non-infected clutch with independent eggs and interstices between them; B) *R. azarai* RG (LGE 7329); C) *R. dorbignyi* RG (unvouchered); D) *R. fernandezae* RG (LGE 7008); E) *R. major* RG (unvouchered); F) *R. arenarum* RM (unvouchered); G) *R. ornata* RC (LGE 8824); H) *R. achalensis* RS (unvouchered); I) *R. spinulosa spinulosa* RS (LGE 6795).

Discussion

Oviposition modes in *Rhinella*

Oocytes of amphibians are enclosed by a thin vitelline membrane, which is composed of glycoproteins that are synthesised and secreted during oocyte maturation in the ovary. Surrounding them are a number of jelly layers produced by oviduct secretions that are deposited around the oocytes as they flow through the different regions of the oviduct (SALTHER 1963, JEGO et al. 1980, ARRANZ et al. 1997). The plesiomorphic and most common oviposition mode reported in *Rhinella* consists of a string, composed of a uni- or bilayered jelly tube with numerous eggs. It is characteristic of the *R. crucifer*, *R. granulosa*, *R. margaritifera*, *R. marina*, and partly of the *R. spinulosa* species groups (see Appendix 1). Eggs within the strings can adopt three well-defined configurations: (i) uniserial, where the eggs are aligned inside smooth (e.g., *R. dorbignyi*, Fig. 2C) or scalloped strings (*R. major* and *R. fernandezae*, Figs. 2D–E); (ii) biserial, where the eggs are staggered inside the tube (e.g., *R. arenarum*, Fig. 2F); or (iii) multiserial, where groups of eggs are laid in a large, thick jelly tube (e.g., *R. achalensis* and *R. spinulosa*; Figs. 2H–I). We consider this latter case to be multiserial strings instead of strands (sensu ALTIG & McDIARMID 2007), because we found structural differences in that the eggs are ordered, and the jelly tube is definitively not flimsy or indistinct but firm and very resistant like that of regular uniserial or biserial strings. *Rhinella achalensis* has so far not been included in phylogenetic analyses, and given that some species of the *R. spinulosa* group deposit their eggs in uniserial strings, multiserial strings could represent a putative synapomorphy of at least an internal clade of this species group. In turn, the open clump of *R. rumbolli* differs remarkably from the strings with different morphologies known in *Rhinella*. Unlike other *Rhinella* eggs, there is no surrounding tube-like matrix enveloping the whole clutch, but the individual capsule of jelly layers is easily distinguished, thick and firm (Appendix 1). Detailed histochemical and molecular studies that unveil the fine structure of the oviducts and their secretions in *R. rumbolli* and other species of *Rhinella* producing strings can help to understand the formation, morphological diversity, and function of the latter.

The “*Rhinella veraguensis*” group, as currently defined, is recovered as polyphyletic in the most inclusive phylogenetic analysis of *Rhinella* (PYRON & WIENS 2011). Part of the *R. veraguensis* group (*R. chavin*, *R. manu*, and *R. nesiotes*) is sister to the *R. acrolopha* group, whereas the remaining species of the *R. veraguensis* group (*R. amboroensis* and *R. veraguensis*) are basal to the latter clade plus the *R. margaritifera* group. The biology of the species of the “*R. veraguensis*” and *R. acrolopha* groups remains poorly known, with no other reports on oviposition other than the one presented here. In several species of these groups, large unpigmented ovarian eggs have been observed (see Appendix 1). The character state unpigmented eggs optimises as a

synapomorphy of the clade composed by *Rhinella chavin*, *R. nesiotes*, *R. manu* (with unknown condition), and species of the *R. acrolopha* group. *Rhinella justiniano*, *R. multiverrucosa*, and *R. yanachaga* also have unpigmented eggs (HARVEY & SMITH 1994, LEHR et al. 2005, 2007) and are most likely related to this clade (for the phylogenetic relationships of *R. yanachaga* see MORAVEC et al. 2014). On the other hand, *R. stanlani* and *R. sternosignata*, two species tentatively assigned to the *R. margaritifera* group, also have unpigmented eggs (LÖTTES & KÖHLER 2000, LA MARCA & MIJARES-URRUTIA 1996). Although both species were not included in a phylogenetic analysis, this condition could be homoplastic considering the position of putative related species of the *R. margaritifera* group.

Several authors have suggested on the basis of the occurrence of large unpigmented oocytes and collecting places distant from water bodies that terrestrial oviposition and an endotrophic developmental mode (direct development or nidicolous larvae) likely occur in the *Rhinella acrolopha* group and in some species of the “*R. veraguensis*” group (DUELLMAN & TOFT 1979, THIBAudeau & ALTIG 1999, LEHR et al. 2007, VAN BOCKLAER et al. 2010, GRANT & BOLIVAR-G. 2014). A redefined *R. acrolopha* group, including the phylogenetically related species of the *R. veraguensis* group (i.e., *R. chavin*, *R. nesiotes*, and *R. yanachaga*; PYRON & WIENS 2011, MORAVEC et al. 2014), would comprise species with large unpigmented eggs and render the *R. veraguensis* group possibly limited to species that breed in lotic water bodies and have exotrophic tadpoles (i.e., *R. chrysophora*, *R. rumbolli*, *R. quechua*, and *R. veraguensis*; McCranie et al. 1989, PRAMUK & LEHR 2005, AGUAYO et al. 2009, HAAD et al. 2014). The close relationship between the two clades provides an opportunity to study evolution of endotrophic development within *Rhinella*. Direct observations on breeding ecology are lacking and still needed, but we certainly could expect more variation in the ovipositional and developmental modes than those currently known in the genus.

Other ovipositional modes reported in Bufonidae, but not noticed or confirmed in *Rhinella*, include jelly bars in *Anaxyrus quercicus* (WRIGHT & WRIGHT 1949; but see the discussion in ALTIG & McDIARMID 2007), rosaries in *Capensibufo rosei* (CHANNING et al. 2012), and terrestrial clumps in *Oreophrynella nigra* (McDIARMID & GORZULA 1989), *Osornophryne guacamayo* (GLUESINKAMP & ACOSTA 2001), and *Xanthophryne tigerinus* (BIJU et al. 2009). Some other species are viviparous, either matrotrophic (*Nimbaphrynoides occidentalis*; ANGEL & LAMOTTE 1944) or lecithotrophic (*Nectophrynoides* spp.; see the revision by LIEDTKE et al. 2014). Descriptions of the breeding biology in poorly-known species of the *R. veraguensis* and *R. acrolopha* groups will contribute with biological data that allow us to understand more thoroughly the diversification in the reproductive biology in *Rhinella* and determine if the shallow clump observed in *R. rumbolli* can have systematic relevance or if it constitutes an autapomorphy of this species.

Clutch infection by Saprolegniaceae

Our report of infection by Saprolegniaceae on eggs of *Rhinella rumbolli* is the first for the genus. Nevertheless, tadpoles of *R. marina* (as *Bufo marinus*) infected with *Aphanomyces* (Saprolegniaceae) were previously reported for two alien populations in Australia (BERGER et al. 2001). Other bufonids from which this infection known to affect their eggs and embryos are *Anaxyrus americanus* (GOMEZ-MESTRE et al. 2006, TOUCHON et al. 2006), *A. boreas* (BLAUSTEIN et al. 1994, KIESECKER et al. 2001), *Bufo bufo* (ROBINSON et al. 2003), and *Epidalea calamita* (BANKS & BEEBEE 1988, FERNÁNDEZ-BENÉITEZ et al. 2008). Further observations coupled with an appropriate experimental design are mandatory to elucidate several aspects related to this fungal infection. First, we are not certain whether the infection is parasitic or a simple colonization of already dead eggs. Epidemiological aspects such as incidence in clutches and eggs, mortality rates, non-lethal effects, etc., should be addressed. Some hints from other bufonids are useful in this regard. Results in *Anaxyrus americanus* reveal that although high percentages of clutches are infected, the mean mortality tends to be low (only 25%), pointing to some defence or palliative mechanism. In some cases, water mould is ingested by larger tadpoles of the same or different species, and the spread of infection throughout the clutch is curbed thus (GOMEZ-MESTRE et al. 2006). Among non-lethal effects, an induced early hatching has been proven in embryos of *A. americanus*, which can bring their hatching forward by up to 36% compared to control embryos (GOMEZ-MESTRE et al. 2006). We observed hatched stage 17–18 embryos in situ, but it is uncertain if this is a regular event in this species, since early hatching is typical of at least some *Rhinella* (e.g., *R. arenarum*: St. 16; DEL CONTE & SIRLIN 1951). In the case of *R. rumbolli*, it should be tested whether the high incidence and rapid growth of water mould on its clutches is related to the absence of a jelly tube enveloping the whole clump.

Acknowledgements

We thank E. O. LAVILLA, S. KRETZSCHMAR, M. CÁNEPA (FML) for allowing us to study vouchers under their care. We are grateful to G. SCROCCHI, J. C. STAZZONELLI, J. and S. BALDO, and Y. ARZAMENDIA for their company, logistics, and help during fieldwork. We also thank D. CARDOZO, J. M. FERRO, J. LESCANO, D. FENOLIO, and M. MENIN for sharing information or photographs. We thank the Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), ANPCyT, Universidad de Buenos Aires, and the Fundação de Amparo a Pesquisa do Estado de São Paulo for their financial support: PIP 1122008010-2422, 112201101-00875, PIP 11220110100889; PICT 2007-2202, 2011-1524, 2011-1895, 2012-2687 and 2013-404; UBACyT 20020090200727 and 20020110200213; and FAPESP procs. 2012/10000-5 and 2013/50741-7.

References

- AGUAYO, R., E. O. LAVILLA, M. F. VERA CANDIOTI & T. CAMACHO (2009): Living in fast-flowing water: morphology of the gastromyzophorous tadpole of the bufonid *Rhinella quechua* (*R. veraguensis* group). – *Journal of Morphology*, **270**: 1431–1442.
- AICHINGER, M. (1992): Fecundity and breeding sites of an anuran community in a seasonal tropical environment. – *Studies on Neotropical Fauna and Environment*, **27**: 9–18.
- ALTIG, R. & R. W. MCDIARMID (2007): Morphological diversity and evolution of egg and clutch structure in amphibians. – *Herpetological Monographs*, **21**: 1–32.
- ANGEL, F. & M. LAMOTTE (1944): Un crapaud vivipare d'Afrique Occidentale. – *Annales des Sciences Naturelles, Zoologie*, **6**: 63–89.
- ARRANZ, S., I. ALBERTALI & M. CABADA (1997): *Bufo arenarum* egg jelly coat: purification and characterization of two highly glycosylated proteins. – *Biochemical Journal*, **323**: 307–312.
- ANSTIS, M. (2013): Tadpoles and frogs of Australia. – London: New Holland.
- BALDISSERA, F. A., U. CARAMASCHI & C. F. B. HADDAD (2004): Review of the *Bufo crucifer* species group, with descriptions of two new related species (Amphibia, Anura, Bufonidae). – *Arquivos do Museu Nacional, Rio de Janeiro*, **62**: 255–282.
- BANKS, B. & T. J. C. BEEBEE (1988): Reproductive success of natterjack toads *Bufo calamita* in two contrasting habitats. – *The Journal of Animal Ecology*, **57**: 475–492.
- BERGER, L., R. SPEARE, A. THOMAS & A. HYATT (2001): Mucocutaneous fungal disease in tadpoles of *Bufo marinus* in Australia. – *Journal of Herpetology*, **35**: 330–335.
- BIJU, S. D., I. VAN BOCXLAER, V. B. GIRI, S. P. LOADER & F. BOSUYT (2009): Two new endemic genera and a new species of toad (Anura: Bufonidae) from the Western Ghats of India. – *BMC Research Notes*, **2**.
- BLAUSTEIN, A. R., D. GRANT HOKIT, R. K. O'HARA & R. A. HOLT (1994): Pathogenic fungus contributes to amphibian losses in the Pacific Northwest. *Biological Conservation*, **67**: 251–254.
- BLOTTO, B. L., M. O. PEREYRA & D. BALDO (2014): The tadpole of *Rhinella azarai* (GALLARDO, 1965) with comments on larval morphology in the *Rhinella granulosa* species group (Anura: Bufonidae). – *Journal of Herpetology*, **48**: 434–438.
- CABRERA, A. L. (1994): Regiones fitogeográficas argentinas. – pp. 1–85 in: KUGLER, W. F. (ed.): *Enciclopedia argentina de agricultura y jardinería*. – Argentina: Acme.
- CALDWELL, J. P. (1991): A new species of toad in the genus *Bufo* from Pará, Brazil, with an unusual breeding site. – *Papéis Avulsos Zoologia* (São Paulo), **37**: 389–400.
- CALDWELL, J. P. & M. C. DE ARAÚJO (2004): Historical and ecological factors influence survivorship in two clades of phytoelm-breeding frogs (Anura: Bufonidae, Dendrobatidae). – pp. 11–21 in: LEHTINEN, R. M. (ed.): *Ecology and evolution of phytoelm-breeding anurans*. – Ann Arbor, Michigan, Miscellaneous Publications Museum of Zoology University of Michigan.
- CALDWELL, J. P. & D. B. SHEPARD (2007): Calling Site Fidelity and Call Structure of a Neotropical Toad, *Rhinella ocellata* (Anura: Bufonidae). – *Journal of Herpetology*, **41**: 6 11–621.

- CARAMASCHI, U. & H. NIEMEYER (2003): Nova espécie do complexo de *Bufo margaritifera* (Laurenti, 1768) do estado do Mato Grosso do Sul, Brasil (Amphibia, Anura, Bufonidae). – Boletim do Museu Nacional, Nova Série, Zoologia, **501**: 1–16.
- CARRIZO, G. R. (1992): Cuatro especies nuevas de anuros (Bufonidae: *Bufo* e Hylidae: *Hyla*) del norte de la Argentina. – Cuadernos de Herpetología, **7**: 14–23.
- CARVALHO E SILVA, A. M. P. T. CARVALHO E SILVA & S. P. CARVALHO E SILVA (1994): Données sur la biologie et description des larves de *Bufo pygmaeus* Myers et Carvalho (Amphibia, Anura, Bufonidae). – Revue Française d'Aquariologie, **21**: 53–56.
- CEL, J. M. (1980): Amphibians of Argentina. – Monitore Zoologico Italiano, N. S., **2**: 1–609.
- CHANNING, A., M.-O. RÖDEL & J. CHANNING (2012): Tadpoles of Africa. The biology and identification of all known tadpoles in sub-Saharan Africa. – Frankfurt am Main, Chimaira Edition.
- CHAPARRO, J. C., J. B. PRAMUK & A. G. GLUESENKAMP (2007): A new species of arboreal *Rhinella* (Anura: Bufonidae) from cloud forest of southeastern Peru. – Herpetologica, **63**: 203–212.
- CRUMP, M. L. (1974): Reproductive strategies in a tropical anuran community. – Miscellaneous Publications of the Museum of Natural History of the University of Kansas, **61**: 1–68.
- DEL CONTE, E. & L. SIRLIN (1951): The first stages of *Bufo arenarum* development. – Acta Zoológica Lilloana, **12**: 495–499.
- DIXO, M. & V. K. VERDADE (2006): Herpetofauna de serrapilheira da Reserva Florestal de Morro Grande, Cotia (SP). – Biota Neotropica, **6**: 1–20.
- DUELLMAN, W. E. (1978): The biology of an equatorial herpetofauna in Amazonian Ecuador. – University of Kansas Publications, Museum of Natural History, **65**: 1–352.
- DUELLMAN, W. E. (2005): Cusco Amazonico. The lives of amphibians and reptiles in an Amazonian rainforest. – Ithaca: Cornell University Press.
- DUELLMAN, W. E. & R. SCHULTE (1992): Description of a new species of *Bufo* from northern Peru with comments on phenetic groups of South American toads (Anura: Bufonidae). – Copeia, **1992**: 162–172.
- DUELLMAN, W. E. & C. A. TOFT (1979): Anurans from the Serranía de Sira, Amazonian Perú: Taxonomy and biogeography. – Herpetologica, **35**: 60–70.
- FENOLIO, D. B., J. R. MENDELSON & W. W. LAMAR (2012): A new diagnosis and description of variation among adult *Rhinella ceratophrys* (Boulenger) (Amphibia: Bufonidae), with notes on ecology and distribution. – South American Journal of Herpetology, **7**: 9–15.
- FERNÁNDEZ, K. (1927): Sobre la biología y reproducción de batracios argentinos. Segunda parte. – Boletín Academia Nacional de Ciencias de Córdoba, **29**: 271–320.
- FERNÁNDEZ-BENÍTEZ, M. J., M. E. ORTIZ-SANTALIESTRA, M. LIZANA & J. DIÉGUEZ-URIBEONDO (2008): *Saprolegnia diclina*: another species responsible for the emergent disease 'Saprolegnia infections' in amphibians. – FEMS Microbiological Letters, **279**: 23–29.
- FROST, D. R. (2014): Amphibian Species of the World: an online reference. Version 6.0 (accessed 5 June 2014). – Electronic Database accessible at <http://research.amnh.org/herpetology/amphibia/index.html>. – American Museum of Natural History, New York, USA.
- GALLARDO, J. M. (1969): La distribución de las subespecies de *Bufo granulosus* Spix: su fidelidad a los sistemas hidrográficos Sudamericanos. – Ciencia e Investigación, **25**: 406–416.
- GLUESENKAMP, A. G. & N. ACOSTA (2001): Sexual dimorphism in *Osornophryne guacamayo* with notes on natural history and reproduction in the species. – Journal of Herpetology, **35**: 148–151.
- GOMEZ-MESTRE, I., J. C. TOUCHON & K. M. WARKETIN (2006): Amphibian embryo and parental defenses and a larval predator reduce egg mortality from water mold. – Ecology, **87**: 2570–2581.
- GRANT, T. & W. BOLIVAR-G (2014): A new species of semi-arboreal toad with a salamander-like ear (Anura: Bufonidae: *Rhinella*). – Herpetologica, **70**: 198–210.
- GUAYARA-BARRAGÁN, M. G. & M. H. BERNAL (2012): Fecundidad y fertilidad en once especies de anuros colombianos con diferentes modos reproductivos. – Caldasia, **34**: 483–496.
- HAAD, M. B., F. VERA CANDIOTI & D. BALDO (2014): The stream tadpoles of *Rhinella rumbolli* (Anura: Bufonidae). – Herpetologica, **70**: 184–197.
- HADDAD, C. F. B. & C. P. A. PRADO (2005): Reproductive Modes in Frogs and Their Unexpected Diversity in the Atlantic Forest of Brazil. – BioScience, **55**: 207–217.
- HARVEY, M. B. & E. N. SMITH (1994): A new species of *Bufo* (Anura: Bufonidae) from cloud forest in Bolivia. – Herpetologica, **50**: 32–38.
- HÖDL, W. (1990): Reproductive diversity in Amazonian lowland frogs. – Fortschritte der Zoologie **38**: 41–60.
- JEGO, P., J. JOLY & C. BOISSEAU (1980): Les gangues ovulaires des amphibiens (protéines sécrétées par l'oviducte) et leurs rôles dans la fécondation. – Reproduction Nutrition Développement, **20**: 557–567.
- JOFRÉ, G. M., C. J. READING & I. E. DI TADA (2005): Breeding behaviour and reproduction in the Pampa de Achala toad, *Bufo achalensis*. – Amphibia-Reptilia, **26**: 451–458.
- KENNY, J. S. (1969): The Amphibia of Trinidad. – Studies on the fauna of Curaçao and other Caribbean islands, **29**: 1–78.
- KIESECKER, J. M., A. R. BLAUSTEIN & L. K. BELDEN (2001): Complex causes of amphibians population declines. – Nature, **410**: 681–684.
- KÖHLER, J. (2000): Amphibian diversity in Bolivia: a study with special reference to montane forest regions. – Bonner Zoologische Monographien, **48**: 1–243.
- LA MARCA, E. & A. MIJARES-URRUTIA (1996): Taxonomy and geographic distribution of a northwestern Venezuelan toad (Anura, Bufonidae, *Bufo sternosignatus*). – Alytes, **14**: 101–114.
- LEHR, E., G. KÖHLER, C. AGUILAR & E. PONCE (2001): New Species of *Bufo* (Anura: Bufonidae) from central Peru. – Copeia, **2001**: 216–223.
- LEHR, E., J. B. PRAMUK & M. LUNDBERG (2005): A new species of *Bufo* (Anura: Bufonidae) from Andean Peru. – Herpetologica, **61**: 308–318.
- LEHR, E., J. B. PRAMUK, S. B. HEDGES & J. H. CÓRDOVA (2007): A new species of arboreal *Rhinella* (Anura: Bufonidae) from Yanachaga-Chemillén National Park in central Peru. – Zootaxa, **1662**: 1–14.

- LIEDTKE, H. C., H. MÜLLER, J. HAFNER, P. NAGEL & S. P. LOADER (2014): Interspecific patterns for egg and clutch sizes of African Bufonidae (Amphibia: Anura). – *Zoologischer Anzeiger*, **253**: 309–315.
- LIMA, A. P., W. E. MAGNUSSEN, M. MENIN, L. K. ERDTMANN, D. J. RODRIGUES, C. KELLER & W. HÖDL (2006): Guide to the frogs of Reserva Adolpho Ducke, Central Amazonia. – Manaus, Attema Design Editorial.
- LÖTTERS, S. & J. KÖHLER (2000): A new toad of the *Bufo typhoni* complex. – *Spixiana*, **23**: 293–303.
- LOURENÇO, A. C. C., D. BAETA, A. C. L. DE ABREU & J. P. POMBAL (2010): Tadpole and advertisement call of *Rhinella pombali* (Baldiessa, Caramaschi & Haddad, 2004) (Amphibia, Anura, Bufonidae). – *Zootaxa*, **2370**: 65–68.
- LYNCH, J. D. & J. M. RENJIFO (1990): Two new toads (Bufonidae: *Rhombophryne*) from the northern Andes of Colombia. – *Journal of Herpetology*, **24**: 364–371.
- MCCRANIE, J. R., L. D. WILSON & K. L. WILLIAM (1989): A new genus and species of toad (Anura: Bufonidae) with an extraordinary stream-adapted tadpole from northern Honduras. – *Journal of Occasional Papers of the Museum of Natural History, the University of Kansas*, **129**: 1–18.
- MCDIARMID, R. W. & S. GORZULA (1989): Aspects of the reproductive ecology and behavior of the tepui toads, genus *Oreophrynella* (Anura, Bufonidae). – *Copeia*, **1989**: 445–451.
- MENIN, M., D. J. RODRIGUES & A. P. LIMA (2006): The tadpole of *Rhinella proboscidea* (Anura: Bufonidae) with notes on adult reproductive behavior. – *Zootaxa*, **1258**: 47–56.
- MIJARES-URRUTIA, A. & A. ARENDS (2001): A new toad of the *Bufo margaritifera* complex (Amphibia: Bufonidae) from Northwestern Venezuela. – *Herpetologica*, **57**: 523–531.
- MORAVEC, J., E. LEHR, J. C. CUSI, J. H. CORDOVA & V. GVOŽDÍK (2014): A new species of the *Rhinella margaritifera* species group (Anura, Bufonidae) from the montane forest of the Selva Central, Peru. – *ZooKeys*, **371**: 35–56.
- NARVAES, P. & M. T. RODRIGUES (2009): Taxonomic revision of *Rhinella granulosa* species group (Amphibia, Anura, Bufonidae), with a description of a new species. – *Arquivos de Zoologia, São Paulo*, **40**: 1–73.
- PEROTTI, G. M. (1994): Aporte preliminares sobre la reproducción en una comunidad de anuros chaqueños en Argentina. – *Cuadernos de Herpetología*, **8**: 39–50.
- PEROTTI, M. G. (1997): Modos reproductivos y variables reproductivas cuantitativas de un ensamble de anuros del Chaco semiárido, Salta, Argentina. – *Revista Chilena de Historia Natural*, **70**: 277–288.
- PRAMUK, J. B. (2006): Phylogeny of South American *Bufo* (Anura: Bufonidae) inferred from combined evidence. – *Zoological Journal of the Linnean Society*, **146**: 407–452.
- PRAMUK, J. B. & F. KADIVAR (2003): A new species of *Bufo* (Anura: Bufonidae) from southern Ecuador. *Herpetologica*, **59**: 270–283.
- PRAMUK, J. B. & E. LEHR (2005): Taxonomic status of *Atelophryniscus chrysophorus* McCranie, Wilson, and Williams, 1989 (Anura: Bufonidae) inferred from phylogeny. – *Journal of Herpetology*, **39**: 610–618.
- PYRON, R. A. & J. J. WIENS (2011): A large-scale phylogeny of Amphibia including over 2800 species, and a revised classification of extant frogs, salamanders, and caecilians. *Molecular Phylogenetics and Evolution*, **61**: 543–583.
- QUIROGA, L. B. & E. A. SANABRIA (2012): Variation in reproductive parameters of *Rhinella arenarum* (Hensel, 1867) (Anura: Bufonidae) between the reproductive and post-reproductive periods. – *Belgian Journal of Zoology*, **142**: 68–73.
- RIVERO, J. A. & C. J. CASTAÑO (1990): A new and peculiar species of *Rhombophryne* (Amphibia: Bufonidae) from Antioquia, Colombia. – *Journal of Herpetology*, **24**: 1–5.
- ROBINSON, J., R. A. GRIFFITHS & P. JEFFRIES (2003): Susceptibility of frog (*Rana temporaria*) and toad (*Bufo bufo*) eggs to invasion by *Saprolegnia*. – *Amphibia-Reptilia*, **24**: 261–268.
- RODRÍGUEZ, L. O. & W. E. DUELLMAN (1994): Guide to the frogs of the Iquitos region, Amazonian Peru. – The University of Kansas Natural History Museum Special Publication, **22**: 1–80.
- SALTHER, S. N. (1963): The egg capsules in the Amphibia. – *Journal of Morphology*, **113**: 161–171.
- SAVAGE, J. M. (2002): The amphibians and reptiles of Costa Rica. – Chicago, The University of Chicago.
- SINSCH, U., I. E. DI TADA & A. L. MARTINO (2001): Longevity, demography and sex-specific growth of the Pampa de Achala toad, *Bufo achalensis* Ceí, 1972. – *Studies on Neotropical Fauna and Environment*, **36**: 95–104.
- SYMPSON, V., F. G. JARA & C. A. ÚBEDA (2006): *Bufo spinulosus papillosus* (NCN), Reproduction. – *Herpetological Review*, **37**: 200–201.
- THIBAudeau, G. & R. ALTIG (1999): Endotrophic anurans. Development and evolution. – pp. 170–188 in: McDIARMID, R. W. & R. ALTIG (eds): The biology of anuran larvae. – Chicago, Chicago University Press.
- TOUCHON, J. C., I. GOMEZ-MESTRE & K. M. WARKENTIN (2006): Hatching plasticity in two temperate anurans: responses to a pathogen and predation cues. – *Canadian Journal of Zoology*, **84**: 556–563.
- TRUEB, L. (1971): Phylogenetic relationships of certain neotropical toads with the description of a new genus (Anura: Bufonidae). – Los Angeles County Museum of Natural History, **216**: 1–40.
- URRA, F. A. (2013): Síntesis del conocimiento actual sobre los sapos *Rhinella atacamensis*, *R. arunco* y *R. spinulosa*. – *La Chiricoca*, **3**: 4–15.
- VAN BOCKLAER, I., S. P. LOADER, K. ROELANTS, S. D. BIJU, M. MENEGON & F. BOSSUYT (2010): Gradual adaptation toward a range-expansion phenotype initiated the global radiation of toads. – *Science*, **327**: 679–682.
- VÉLEZ-R., C. M. & P. M. RUIZ-C. (2002): A new species of *Bufo* (Anura: Bufonidae) from Colombia. – *Herpetologica*, **58**: 453–462.
- WRIGHT, A. H. & A. A. WRIGHT (1949): Handbook of frogs and toads of the United States and Canada. – Ithaca, NY: Comstock.
- YANOSKY, A. A., C. MERCOLLI & J. A. DIXON (1993): Field ecology of the pygmy toad *Bufo pygmaeus* (Anura: Bufonidae), in northeastern Argentina with notes on sympatric sibling species of the *granulosus* group. – *Bulletin of the Maryland Herpetological Society*, **33**: 66–77.

Appendix 1

Variation in reproductive traits of *Rhinella* spp. All data obtained from clutches, except those indicated with a (from mature ovarian oocytes) or b (clutch and mature ovarian oocytes). * – Data inferred from photographs in publications. Measurements are in mm.

	Oviposition mode	Number of eggs	Ovum pigmentation	Ovum size	String / Strand diameter
<i>Rhinella acrolopha</i> group					
<i>R. acrolopha</i> ^a	?	290 (145 in one side) ¹	unpigmented ¹	2.5 ¹	?
<i>R. festae</i> ^a	?	“few” ¹	unpigmented ¹	1.8 ¹	?
<i>R. lindae</i> ^a	?	7 + 11 ²	unpigmented ²	“large” ²	?
<i>R. macrorrhina</i> ^a	?	“few” ¹	unpigmented ¹	2 ¹	?
<i>R. nicefori</i> ^a	?	“few” ¹	unpigmented ¹	1.8 ¹	?
<i>R. paraguas</i> ^a	?	61 ³	unpigmented ³	2.24±0.23 ³	?
<i>R. tenrec</i> ^a	?	?	unpigmented ⁴	2.5–2.7 ⁴	?
<i>R. crucifer</i> group					
<i>R. ornata</i>	strings (uniserial) ^{5,6}	>831 (N=1) (incomplete clutch) ⁶	animal pole pigmented ⁶	1.84±0.02 ⁶	3.70±0.25 ⁶
<i>R. pombali</i>	strings (biserial) ^{7*}	9952 ⁷	completely pigmented ⁷	?	?
<i>R. granulosa</i> group					
<i>R. azarai</i>	strings (uniserial, scalloped) ^{6,8}	3770–7548 ^{6,8}	animal pole pigmented ^{6,8}	1.19±0.06 (1.06–1.33) ⁶	1.72±0.19 (1.48–1.97) ⁶
<i>R. bergi</i> (as <i>Bufo pygmaeus</i>)	strings ⁹	?	?	1.00 ⁹	1.80 ⁹
<i>R. dorbignyi</i>	strings ¹⁰	?	?	?	?
	strings ^{10,11}	9000 ¹¹	?	?	3.00–3.50 ¹¹
<i>R. fernandezae</i> (as <i>B. dorbignyi</i> in ¹¹)	strings (biserial) ⁶	1497 ⁶	animal pole pigmented ⁶	1.13±0.06 (1.02–1.25) 1.29±0.04 (with vitelline membrane: 1.52±0.09) ⁶	2.87±0.14 ⁶
<i>R. merianae</i> (as <i>B. granulatus</i> in ¹²)	strings ¹²	983 ¹²	?	<1.0 ¹²	?
	strings ¹³	900 ¹³	?	?	?
<i>R. humboldti</i> (as <i>B. g. beebei</i> in ¹⁴ and <i>R. granulosa</i> in ¹⁵)	strings ¹⁴	1335–4391 ¹⁵	?	1.37±0.10 (1.23–1.55) ¹⁵	?
<i>R. major</i> ^a (as <i>B. granulatus</i> in ¹⁶ and ¹⁷)	?	7784 (1901–13195) ^{16,17}	pigmented ^{16,17}	0.52 ^{16,17}	?
	strings (uniserial, scalloped) ⁶	6269 (N=1) ⁶	animal pole pigmented ⁶	0.95±0.07 (0.84–1.07) ⁶	1.35±0.03 (1.3–1.39) ⁶
<i>R. pygmaea</i>	strings (uniserial) ¹⁸	4140 ¹⁸	pigmented ^{18*}	1.20 (with vitelline membrane: 1.40) ¹⁸	1.60 ¹⁸
<i>R. margaritifera</i> group					
<i>R. castaneotica</i>	strings (uniserial) ^{19*}	61–387 ^{19,20}	pigmented ^{19*}	?	?
<i>R. ceratophrys</i> ^a	?	1000–1500 ²¹	pigmented ²¹	<1.00 ²¹	?
<i>R. cristinae</i> ^a	?	“hundreds” ²²	animal pole pigmented ²²	1.20–1.60 ²²	?
	?	310–2500 ²³	animal pole pigmented ²³	<2.00 ²³	?
	strings ²⁴	?	pigmented ²⁴	“small” ²⁴	?
<i>R. margaritifera</i> (as <i>B. typhonius</i> in ^{23–26})	strings ²⁵	1150–2431 ²⁵	?	1.29 ²⁵	?
	strings ²⁵	1226 ²⁵	?	1.21 ²⁵	?
	strings ²⁶	765–2500 ²⁶	pigmented ²⁶	?	?
	strings ²⁷	1165–1630 ²⁷	animal pole pigmented ²⁷	1.30–1.50 ²⁷	?
<i>R. ocellata</i>	strings (uniserial) ²⁸	2240 ²⁸	pigmented ²⁸	1.70–1.79 ²⁸	?
	?	450 ¹³	?	?	?
<i>R. proboscidea</i>	strings (uniserial) ²⁹	439–473 ²⁹	animal pole pigmented ²⁹	2.76 (2.4–3.4) ²⁹ (with capsule: 3.7–4.7) ³⁰	?
<i>R. scitula</i>	strings ³¹	?	?	?	?
<i>R. sclerocephala</i> ^a	?	“hundreds, maybe thousand” ³²	animal pole pigmented ³²	1.40–1.80 ³²	?
<i>R. stanlani</i>	?	?	unpigmented ³³	?	?

Oviposition modes in *Rhinella*

	Oviposition mode	Number of eggs	Ovum pigmentation	Ovum size	String / Strand diameter
<i>R. sternosignata</i>	?	?	unpigmented ³⁴	1.4–1.9 ³⁴	?
<i>R. yunga</i>	?	“numerous” ³⁵	pigmented ³⁵	“small” ³⁵	?
<i>R. marina</i> group					
	strings (biserial) ¹¹	40000 ¹¹	?	?	4.00–5.00 ¹¹
	strings ³⁶	4000–5000 ³⁶	pigmented ³⁶	?	?
<i>R. arenarum</i>	strings ³⁷	33219 ^{37,a}	animal pole pigmented ^{37,a}	1.06±0.09 ^{37,a}	?
	strings (biserial) ⁶	?	animal pole pigmented ⁶	1.23±0.22 (with vitelline membrane: 1.40±0.05) ⁶	3.15±0.17 ⁶
<i>R. cerradensis</i>	strings (biserial) ⁶	?	animal pole pigmented ⁶	2.09±0.08 ⁶	6.44±0.08 ⁶
<i>R. icterica</i>	strings ⁵	?	?	?	?
	?	4240–12700 ^{23,a}	animal pole pigmented ^{23,a}	<1.5 ^{23,a}	?
	strings ¹²	8598 ¹²	?	<1.5 ¹²	?
	strings ³⁸	5000–25000 ³⁸	animal pole pigmented ³⁸	1.7–2.0 ³⁸	?
<i>R. marina</i>	strings ¹³	4000–10000 ¹³	?	?	?
	strings ¹⁵	10000–20000 ^{15,b}	?	1.50±0.03 (1.46–1.54) ^{15,b}	?
	strings ³⁹	7000–35000 ³⁹	animal pole pigmented ³⁹	1.5–1.6 ³⁹	?
<i>R. poeppigii</i>	strings ⁴⁰	?	?	?	?
<i>R. schneideri</i> (as <i>B. paracnemis</i>)	strings ³⁶	?	pigmented ³⁶	1.8 ³⁶	?
<i>R. spinulosa</i> group					
	strings (multiserial) ⁴¹	700–2548 ^{41,b}	?	?	?
<i>R. achalensis</i>	strings (multiserial) ^{42*}	?	pigmented ^{42*}	?	?
	strings (multiserial) ⁶	?	pigmented ⁶	?	?
<i>R. amabilis</i> ^a	?	1436 ⁴³	pigmented ⁴³	0.47 (0.36–0.52) ⁴³	?
	strings ^{44*}	?	pigmented ⁴⁴	?	?
<i>R. arunco</i>	strings (biserial) ⁴⁵	?	pigmented ⁴⁵	?	?
<i>R. atacamensis</i>	strings ⁴⁴	?	pigmented ⁴⁴	?	?
	strings (multiserial) ⁴⁴	“thousands” ⁴⁴	pigmented ⁴⁴	?	?
<i>R. spinulosa</i> (as <i>B. spinulosus papillosus</i> in ⁴⁵)	strings ¹¹	?	pigmented ¹¹	1.3 (cf.) ¹¹	5.00 ¹¹
	strings (multiserial) ⁴⁶	3250–6400 ⁴⁶	animal pole pigmented ⁴⁶	1.56±0.09 ⁴⁶	>8.30 ⁴⁶
	strings (multiserial) ⁶	>1401 (incomplete) ⁶	animal pole pigmented ⁶	1.64±0.06 ⁶	5.9±0.18 ⁶
“<i>R. veraguensis</i>” group					
<i>R. chavin</i> ^a	?	266–286 ⁴⁷	unpigmented ⁴⁷	1.82–3.05 ⁴⁷	?
<i>R. fissipes</i> ^a	?	?	pigmented ⁴⁰	~1 ⁴⁰	?
<i>R. justinianoi</i> ^a	?	?	unpigmented ⁴⁸	?	?
<i>R. manu</i> ^a	?	35–40 ⁴⁹	?	2.00 ⁴⁹	?
<i>R. multiverrucosa</i> ^a	?	234–290 ⁵⁰	unpigmented ⁵⁰	2.12–2.30 ⁵⁰	?
<i>R. nesiotes</i> ^a	?	20 ⁵¹	unpigmented ⁵¹	1.2 ⁵¹	?
<i>R. quechua</i> ^a	?	1014 ⁶	animal pole pigmented ⁶	1.27±0.05 (1.19–1.37) ⁶	?
<i>R. rumbolli</i>	shallow clump ⁶	2177 ⁶	animal pole pigmented ⁶	1.91±0.09 (with capsule: 3.85±0.36) ⁶	–
<i>R. yanachaga</i> ^a	?	136 ⁵²	unpigmented ⁵²	?	?

References. ¹TRUEB 1971, ²RIVERO & CASTAÑO 1990, ³GRANT & BOLIVAR-G. 2014, ⁴LYNCH & RENJIFO 1990, ⁵DIXO & VERDADE 2006, ⁶present work, ⁷LOURENÇO et al. 2010, ⁸BLOTTO et al. 2014, ⁹YANOSKY et al. 1993, ¹⁰GALLARDO 1969, ¹¹FERNÁNDEZ 1927, ¹²HÖDL 1990, ¹³LIMA et al. 2006, ¹⁴KENNY 1969, ¹⁵GUAYARA-BARRAGÁN & BERNAL 2012, ¹⁶PEROTTI 1994, ¹⁷PEROTTI 1997, ¹⁸CARVALHO E SILVA & CARVALHO E SILVA 1994, ¹⁹CALDWELL & DE ARAÚJO 2004, ²⁰CALDWELL 1991, ²¹FENOLIO et al. 2012, ²²VÉLEZ-R. & RUIZ-C. 2002, ²³CRUMP 1974, ²⁴DUELLMAN 1978, ²⁵AICHINGER 1992, ²⁶RODRÍGUEZ & DUELLMAN 1994, ²⁷DUELLMAN 2005, ²⁸CALDWELL & SHEPARD 2007, ²⁹MENIN et al. 2006, ³⁰M. MENIN pers. comm., ³¹CARAMASCHI & NIEMEYER 2003, ³²MIJARES-URRUTIA & ARENDS 2001, ³³LÖTTERS & KÖHLER 2000, ³⁴LA MARCA & MIJARES-URRUTIA 1996, ³⁵MORAVEC et al. 2014, ³⁶CEI 1980, ³⁷QUIROGA & SANABRIA 2012, ³⁸SAVAGE 2002, ³⁹ANSTIS 2013, ⁴⁰KÖHLER 2000, ⁴¹JOFRÉ et al. 2005, ⁴²SINSCH et al. 2001, ⁴³PRAMUK & KADIVAR 2003, ⁴⁴URRA 2013, ⁴⁵D. FENOLIO pers. comm., ⁴⁶SYMPSON et al. 2006, ⁴⁷LEHR et al. 2001, ⁴⁸HARVEY & SMITH 1994, ⁴⁹CHAPARRO et al. 2007, ⁵⁰LEHR et al. 2005, ⁵¹DUELLMAN & TOFT 1979, ⁵²LEHR et al. 2007. Species referenced in ^{1, 2, 4} as *Rhamphophryne* and ^{5, 9–12, 14, 16–20, 22–27, 31–34, 36, 38, 40–43, 46–48, 50–51} as *Bufo*.

Appendix 2

Examined material

Rhinella azarai – Argentina: Misiones: Departamento Capital: Garupá, Barrio Santa Helena; LGE 3821 (incomplete clutch); LGE 5822 (complete clutch with 5043 eggs); LGE 3736 (complete clutch with 3770 eggs, BLOTTO et al. 2014); LGE 3737 (complete clutch with 7548 eggs, BLOTTO et al. 2014); LGE 3738 (complete clutch with 4053 eggs); Departamento Concepción: Concepción de la Sierra; LGE 7329 (incomplete clutch).

Rhinella cf. cerradensis – Argentina: Corrientes: Departamento Santo Tomé: Gobernador Virasoro; LGE 6338 (clutches from two different pairs); LGE 6339 (incomplete clutch).

Rhinella fernandezae – Argentina: Corrientes: Departamento Ituzaingó: Ituzaingó; LGE 7008 (incomplete clutch); Entre Ríos: Departamento Islas del Ibicuy: Antigua Ruta Nacional N° 12; MACN 39247 (female with ovarian oocytes);

Rhinella major – Argentina: Formosa: Departamento Bermejo: Laguna Yema; LGE 7977, 7982 (two incomplete clutches); LGE 8382 (complete clutch with 6269 eggs).

Rhinella ornata – Argentina: Misiones: Departamento San Pedro: Colonia Victoria, Reserva Freaza; LGE 8824 (incomplete clutch).

Rhinella quechua – Bolivia: Carrasco: Departamento Cochabamba; MACN 46656 (female with ovarian oocytes).

Rhinella rumbolli – Argentina: Jujuy: Departamento Ocloyas: Río Trigo Pampa; LGE 6415 (complete clutch obtained in the laboratory); LGE 6455, 6461–2, 6471, 6475 (five incomplete clutches infected by Saprolegniaceae).

Rhinella spinulosa spinulosa – Argentina: Jujuy: Departamento Tumbaya: El Angosto, 6 kilometers from El Moreno; LGE 8825 (incomplete clutch); Catamarca: Departamento Andalgalá: Laguna del Río Blanco, 3 km. from Río Candado. FML 07171 (incomplete clutch).